arXiv:1209.5227v1 [astro-ph.HE] 24 Sep 2012

Non-stationary magnetic microstructures in stellar thin accretion discs

Giovanni Montani[1, 2, *] and Jacopo Petitta[2, †]
[1] *ENEA - C.R. Frascati, U.T. Fus. (FUSMAG Lab) – Via Enrico Fermi 45, (00044) Frascati (RM), Italy*
[2] *Physics Department, 'Sapienza' University of Rome – Piazzale Aldo Moro 5, (00185) Roma, Italy*

(Dated: September 21, 2012 – Accepted for reviewing by MNRAS)

We examine the morphology of magnetic structures in thin plasma accretion discs, generalizing a stationary ideal MHD model to the time-dependent visco-resistive case. Our analysis deals with small scale perturbations to a central dipole-like magnetic field, giving rise to the periodic modulation of magnetic flux surfaces along the radial direction, which corresponds to the formation of a sequence of toroidal current channels. These microstructures exhibit an exponential damping in time because of the non-zero resistivity coefficient, allowing us to define a lifetime of the configuration which mainly depends on the midplane temperature and on the length scale of the structure itself. If we try to identify a suitable phenomenological scenario where this framework could be applied, it turns out that the steady-state assumption adopted in literature is inconsistent with the evolution time-scale of typical stellar accretion disc systems; we can instead aim to describe peculiar transient events whose duration ranges from months or years.

Keywords: MHD – plasmas – accretion, accretion discs – stars: magnetic field

I. INTRODUCTION

A new perspective on the equilibrium morphology of thin accretion discs has been introduced in [1, 2], where it is outlined how the effect of the plasma back-reaction on the magnetic field of the central object can induce a radially oscillating crystal profile in the magnetic surfaces (see also [3, 4]), eventually breaking up the disc into a ring series, in the limit of a strong non-linear back-reaction. This issue is by its starting point different from the established picture of disc dynamics and its well-known open questions, like the problem of turbulence-enhanced viscosity (for review purposes, you can see the works of [5] and [6]).

The brand new point of view stems from the implementation of an ideal MHD scheme where the Ferraro Corotation Theorem [7] holds, allowing the angular velocity of the disc to be expressed only in terms of the magnetic flux surfaces (for a stationary visco-resistive extension of this model see [8, 9]). In fact, the background centrifugal force is balanced by the central object gravity – resulting in the Keplerian rotation of the disc – as well as by the background magnetic field, since the initial magnetosphere has a current-free morphology. When the back-reaction corrections to the centrifugal force and the Lorentz force are expressed in terms of the magnetic flux function, their balance results in the formation of toroidal current channels embedded in a radial oscillating magnetic structure. A remarkable assumption in constructing this scenario is the pure rotation of the disc, which is not endowed with any poloidal component of the velocity field. Indeed, in many real astrophysical systems, the disc plasma is not very far from the quasi-ideal behaviour, but three main points are in order in this respect:

(i) Given the mean thermodynamic parameters of the plasma in an accreting disc (temperature and number density), the kinetic theory uniquely determines finite non-zero values of the viscosity and resistivity coefficients. Even when the value of such coefficients is rather small, the impact of their existence – i.e. the damping they can induce on the crystal structure – could be not negligible, due to the long lifetime of the accreting systems.

(ii) The Shakura model for accretion within a thin disc configuration [10, 11] relies on very large values of the viscosity coefficient, able to balance the angular momentum transport responsible for a significant non-zero accretion rate. Despite the crystal profile and the Shakura model of accretion are scenarios not directly comparable (e.g. in the crystal profile the purely rotating disc is unable to directly accrete mass), it is of significant interest to understand if the periodic radial profile of the disc still survives in the presence of the visco-resistive effects required to account for the behaviour of a turbulent accreting plasma.

(iii) A non-zero resistivity coefficient is always present in every numerical simulation, at least to reproduce the effect of numerical dissipation due to machine's finite precision [12]. None the less, works like [13] include non-ideal terms to explore the coupling of disc and jet physics, aiming at the problem of angular momentum transport.

In this work we include visco-resistive effects to the crystal profile, allowing for a time dependence of the magnetic surfaces, needed to preserve the consistence of the Generalized Ohm Law. In fact, as far as we include a

---

[*] giovanni.montani@frascati.enea.it

[†] petitta.jacopo@gmail.com

resistive contribution, the azimuthal component of the Ohm Law acquires a term proportional to the toroidal current: this would turn out to be unbalanced in a purely rotating configuration, unless the non-stationarity of the model ensures the presence of a non-zero azimuthal electric field. In such a non-stationary case, we are able to recover the Corotation Theorem and to provide the full consistence of the equilibrium configurations, using assumptions rather natural in the thin disc limit and requiring the toroidal component of the magnetic field to be negligible (we adopt a dipole configuration to characterize the magnetic properties of the central object).

Focusing on the magnetic back-reaction, but neglecting its effect on the mass density distribution (note that this is the most general regime for a non-stationary purely rotating disc), we obtain a crystal profile for the disc magnetic field, isomorphic to the one studied in [1], but for the presence of an exponential damping in time. The characteristic lifetime of this crystal structure depends on the resistivity (or equivalently on the viscosity, since the Prandtl number is constrained to be unity by the consistence of the configuration equations) and crucially on the spatial scale of the radial oscillations. Clearly, when such a lifetime parameter is greater than the average life of the considered accreting system, the resulting profile is mainly stationary and the analysis in [1] is recovered. However, a detailed study of the range of values of the damping lifetime leads to conclude that the crystal profile can not concern the stationary state of a stellar system, because the magnetic microstructures vanish on a timescale much less then the typical lifetime of the accretion disc system.

None the less, the obtained time-dependent scenario has the proper time-scale to account for cataclysmic phenomena. Their duration can be easily fitted in the range of density and temperature which are typical for stellar accretion discs, and in correspondence to radial scales that preserve the request to deal with a local model around a fixed value of the radius (i.e. the length scale of the oscillations is much below the disc radial size). The present analysis is therefore crucial in focusing the correct phenomenological scenario to which the crystal paradigm must be referred to in a stellar system, ruling out this morphology of the magnetic field from the steady state of an accreting stellar configuration.

II. RELEVANT EQUATIONS

Assuming electron pressure gradient is still negligible, the Generalized Ohm's Law retains its validity in the same form of stationary MHD:

$$\vec{E}+\frac{1}{c}\left(\vec{v}\times\vec{B}\right)=\eta\vec{J}\,, \tag{1}$$

obviously meaningful at every different instant t. An important difference stands instead for the expression of electric field descending from Faraday's Law:

$$\vec{E}=-\vec{\nabla}\Phi-\frac{1}{c}\partial_t\vec{A}\,, \tag{2}$$

written in terms of electrostatic potential Φ and vectorial magnetic potential $\vec{A}$. In particular, since the system is axially symmetric, we can say

$$\vec{A}=\frac{\psi}{r}\hat{\mathbf{e}}_\phi \implies \vec{B}=\frac{\vec{\nabla}\psi}{r}\times\hat{\mathbf{e}}_\phi \tag{3}$$

where $\psi\left(r,z^2;t\right)$ is the magnetic flux function, leading to strictly poloidal components of the magnetic field $\vec{B}$.

Since we are dealing with a purely rotating configuration in local approximation (in a narrow annulus around a fixed radius), the azimuthal component of Generalized Ohm's Law (1) becomes:

$$\frac{\partial\psi}{\partial t}-\frac{\eta c^2}{4\pi}\left(\frac{\partial^2\psi}{\partial r^2}+\frac{\partial^2\psi}{\partial z^2}\right)=0\,. \tag{4}$$

Such an expression must be matched with the Induction Equation, namely

$$\frac{\partial\vec{B}}{\partial t}=-\vec{\nabla}\times\left(\vec{v}\times\vec{B}-\frac{c^2\eta}{4\pi}\vec{\nabla}\times\vec{B}\right)\,, \tag{5}$$

and this can be done only if the resistivity it's nearly a constant, precisely if

$$\left|\frac{\vec{\nabla}\eta}{\eta}\right|\ll\left|\frac{\vec{\nabla}(\Delta\psi)}{\Delta\psi}\right|\,, \tag{6}$$

where $\Delta(\cdot)$ is the Laplace operator. This condition ensure that radial and vertical components of Equation (5) give again Equation (4), so they are redundant, whereas the azimuthal one states the following constraint:

$$\vec{\nabla}\omega\times\vec{\nabla}\psi=0\,, \tag{7}$$

telling us that the angular velocity is a flux function $\omega(r,z;t)=\omega(\psi)$. This result generalizes the Corotation Theorem of [7], still holding in the visco-resistive framework provided there is no azimuthal component of the central magnetic field.

Finally, we deal with the MHD momentum conservation equation:

$$\begin{aligned}\rho\left(\partial_t\vec{v}+\left(\vec{v}\cdot\vec{\nabla}\right)\vec{v}\right)=&-\vec{\nabla}p-\rho\vec{\nabla}\chi+\vec{F}_L+\\&+\nu\left[\nabla^2\vec{v}+\frac{1}{3}\vec{\nabla}\left(\vec{\nabla}\cdot\vec{v}\right)\right]\,,\end{aligned} \tag{8}$$

where p is thermodynamic pressure, χ is the gravitational potential of the central object, $\vec{F}_L$ is the Lorentz force, and the viscosity ν is assumed to be a constant. Its azimuthal component gives us the evolution law for the angular velocity:

$$\frac{\partial\omega}{\partial t}-\frac{\nu}{\rho}\left(\frac{\partial^2\omega}{\partial r^2}+\frac{\partial^2\omega}{\partial z^2}\right)=0\,, \tag{9}$$

while the other components retain their stationary forms because of the assumption of pure rotation.

III. PERTURBATIVE SCHEME

Now we can separate the contribution of the background dipole-like magnetic field from the back-reaction induced by the plasma current, i.e.

$$\psi\big(r,z^2;t\big)=\psi_0\big(R_0,z^2\big)+\psi_1\big(r-R_0,z^2;t\big)\,, \tag{10}$$

where R_0 is the fiducial value for our local study, centred on $|r-R_0|\ll R_0$. We highlight that background ψ_0 is a stationary vacuum solution of Laplace's equation (i.e. $\Delta\psi_0=0$), so Equation (4) can be written by means of back-reaction ψ_1 only.

Correspondingly, since Equation (7) constrains the angular velocity to depend upon the magnetic surfaces (and mainly on the back-ground one), an expression for the local study around R_0 is the following:

$$\omega(R_0,r-R_0,z;t)\equiv\omega(\psi)\simeq\omega(\psi_0)+\omega_0'\psi_1\,, \tag{11}$$

with

$$\omega(\psi_0)=\sqrt{\frac{GM_*}{R_0^3}}\,,\quad \omega_0'=\left.\frac{\mathrm{d}\omega}{\mathrm{d}\psi}\right|_{\psi_0}=\text{const.}\,. \tag{12}$$

Note now that the evolution of angular velocity is determined by Equation (9): using the expansion (11), it reduces itself exactly to Equation (4) if

$$\frac{\nu}{\rho}=\frac{c^2\eta}{4\pi}\,, \tag{13}$$

which states that the Magnetic Prandtl Number is of order unity, condition also adopted in [8] with the purpose of recovering the existence of a solution consistent with the Corotation Theorem.

Furthermore, Equation (13) tells us that the height-dependence of $\eta=\eta(\rho)$ is determined only by the density profile, giving a deeper meaning to Equation (6). This can be explained after the introduction of dimensionless variables defined as follows:

$$x=k(r-R_0)\,,\quad u=\frac{z}{\sqrt{\varepsilon_z}H_0},\quad \theta=\frac{t}{\tau}=\frac{k^2\nu}{\rho_0}t\,, \tag{14}$$

and dimensionless functions written as

$$Y\big(x,u^2;\theta\big)=\frac{k\psi_1}{\partial_r\psi_0|_0},\qquad D\big(u^2\big)=\frac{\rho}{\rho_0}\,, \tag{15}$$

where $\rho_0=\rho(x=0,u=0)$ is the density value on the equatorial plane at the fixed radius, H_0 is the half-thickness of the disc, and k^{-1} is the radial scale of the back-reaction flux function ψ_1. The parameter ε_z used in Equation (14) to scale the height-dependence of magnetic surfaces is related to the plasma β-parameter by the relation

$$\beta=8\pi\frac{p_0}{B_{z0}^2}=\frac{1}{3\varepsilon_z^2}\,, \tag{16}$$

with p_0 the back-ground thermodynamic pressure. We can then write down the components of Equation (6) to obtain

$$\frac{\partial_z\eta}{\eta}\simeq\frac{1}{H_0}\ll\frac{1}{\sqrt{\varepsilon_z}H_0}\simeq\frac{\partial_z\psi_1}{\psi_1} \tag{17a}$$

$$\frac{\partial_r\eta}{\eta}\simeq\frac{1}{R_0}\ll k\simeq\frac{\partial_r\psi_1}{\psi_1}\,, \tag{17b}$$

which are identically satisfied if

$$\varepsilon_z\ll 1\,, \tag{18}$$

since Equation (17b) can be restated as

$$kR_0=\frac{R_0}{\varepsilon_z H_0}\gg 1\,, \tag{19}$$

and this is assured for a thin disc. Then it follows that Equation (18) specifies the regime of validity for our treatment.

In terms of these variables and parameters, Equation (4) becomes

$$D\big(u^2\big)\frac{\partial Y}{\partial\theta}-\left(\frac{\partial^2Y}{\partial x^2}+\varepsilon_z\frac{\partial^2Y}{\partial u^2}\right)=0\,, \tag{20}$$

which is a linear parabolic PDE with infinite many solutions. In particular, it admits a separable solution analogue to those found in [2], currently endowed with an explicit time-dependence:

$$Y\big(x,u^2;\theta\big)=A\,F\big(u^2\big)\sin(ax)\,e^{-\theta}\,, \tag{21}$$

with A and a some real constants, whereas the height-dependence is fixed by

$$\varepsilon_z\frac{\mathrm{d}^2F\big(u^2\big)}{\mathrm{d}u^2}-\big(a^2-D\big(u^2\big)\big)\,F\big(u^2\big)=0\,, \tag{22}$$

meaning that it is ultimately fixed after the choice of an equation of state, needed to set an expression for $D\big(u^2\big)$. If we choose a polytropic form for the gravothermal background, such that:

$$p\big(u^2\big)=p_0\,D\big(u^2\big)^{1+1/\Gamma} \tag{23a}$$

$$D\big(u^2\big)=\left[1-\left(\frac{z}{H_0}\right)^2\right]^\Gamma\simeq 1-\Gamma\varepsilon_z u^2\,, \tag{23b}$$



FIG. 1. Behaviour of the back-reaction magnetic surfaces in Equation (24) against dimensionless vertical coordinate, normalized to the gaussian equatorial value obtained at $u = 0$ with $a = a_{\min}$. We fix $\varepsilon_z = 0.05$ and $\Gamma = 1$, then we compute a in the range $[a_{\min}, 1.2]$. As a increases, the plots are drawn as follows: thick, thin, dashed, dotted, dot-dashed. It is worth noting that the gaussian profile has the greater FWHM, while it hasn't the greater initial amplitude.

where Γ is the polytropic index, then we can find an analytic solution of Equation (22) by means of Generalized Hermite Polynomials:

$$F\big(u^2\big) = e^{-\sqrt{\Gamma}u^2/2}\,\mathcal{H}\left[-\frac{\varepsilon_z\sqrt{\Gamma}+a^2-1}{2\varepsilon_z\sqrt{\Gamma}}, \sqrt{\sqrt{\Gamma}u^2}\right], \tag{24}$$

having omitted a constant factor and adopted the notation $\mathcal{H}[\xi, x]$ for the Hermite Polynomial of order ξ in the variable x. Note that these Generalized Polynomials are monotonically decreasing provided their order is negative. Choosing the radial wave-number as $a^2 = a^2_{\min} = 1 - \varepsilon_z\sqrt{\Gamma}$, the solution (24) reproduces the gaussian profile already obtained in [1]. This is the lower limit for the range of values of the parameter, since for $a < a_{\min}$ the solution becomes non-physical; an upper limit doesn't exist instead, although the amplitude of the structure becomes negligible as a increases (see Fig. (1)). Therefore, only a narrow range of values around the unity is suitable to tune the radial wavenumber.

IV. MATCHING WITH PREVIOUS MODELS

We underline that, under the assumption of pure rotation $\vec{v} = \omega R_0 \hat{\mathbf{e}}_\phi$, the only equations explicitly depending on time are those shown in Section II, although there are other interesting equations. Continuity equation in the form:

$$\frac{\partial\rho}{\partial t} + \vec{\nabla}\cdot(\rho\vec{v}) = 0\,, \tag{25}$$

tells us that the density has to keep its steady-state profile, which assures function $D\big(u^2\big)$ doesn't depend on time.

Radial and vertical components of momentum conservation (8) give their stationary formulation too, returning exactly the same dimensionless equations offered in [1, 2], now coupled to Equation (20). This equation introduces the new variable θ, but acts as a closure condition for the system which determines the equilibrium of perturbative pressure $\hat{P}$ and density $\hat{D}$, namely:

$$\partial_{u^2}\hat{P} + \varepsilon_z\hat{D} + 2\left(\partial^2_x Y + \varepsilon_z\partial^2_u Y\right)\partial_{u^2}Y = 0 \tag{26a}$$

$$\begin{aligned}\frac{1}{2}\partial_x\hat{P} + \Big(D\big(u^2\big) + 3\varepsilon_z^2\hat{D}\Big)\,Y +&\\ + \left(\partial^2_x Y + \varepsilon_z\partial^2_u Y\right)(1+\partial_x Y) = 0\,,&\end{aligned} \tag{26b}$$

In the linear regime $Y \ll 1$, when pressure and density perturbations are assumed to be negligible, system (26) reduces itself only to

$$\left(\partial^2_x Y + \varepsilon_z\partial^2_u Y\right) = -D\big(u^2\big)\,Y\,, \tag{27}$$

which is consistent with separable solution (21) and owns the same magnetic structure (although damped in time) developed in [3] or [9], according to the additional approximations considered in those papers. It is interesting to note that Equation (20) could be a closure condition also in the general non-linear regime, but this is forbidden by Equation (25): this suggests that a complete model with the disc decomposition in a ring-like structure resembling the one in [2] (which needs the density to be time-dependent) should exhibit poloidal velocities.

Another solution can be found in the special 'crystal regime' specified by

$$Y \gtrsim 1\,, \quad \hat{D} \ll D\,, \tag{28}$$

occurring when magnetic perturbation are relevant (i.e. back-reaction field is strong) whereas density perturbations are negligible. In this case the system (26) reduces itself to:

$$D\big(u^2\big)\,\partial_\theta Y - \Delta_\varepsilon Y = 0 \tag{29a}$$

$$\partial_{u^2}\hat{P} + 2\Delta_\varepsilon Y\partial_{u^2}Y = 0 \tag{29b}$$

$$\frac{1}{2}\partial_x\hat{P} + D\big(u^2\big)\,Y + \Delta_\varepsilon Y(1+\partial_x Y) = 0\,, \tag{29c}$$

which admits the solution (21) and gives a separable form to pressure too, explicitly:

$$\hat{P}\big(x, u^2; \theta\big) = A^2\,D\big(u^2\big)\,F^2\big(u^2\big)\sin^2(ax)\,e^{-2\theta}\,, \tag{30}$$

where $D\big(u^2\big)$ and $F\big(u^2\big)$ can be expressed via Equation (23b) and Equation (24), respectively.



FIG. 2. Plot of microstructures' lifetime against their length scale via Equation (31) for $T_e = 10^3\,$K (dashed), $10^5\,$K (dotted), $10^7\,$K (dot-dashed), $10^9\,$K (thin line). Shaded region highlights lifetime values whose order of magnitude is from month to year. Thick horizontal line marks the value of 10^7 years. See text for details.

V. MEANING OF DAMPING TIME

It turns out that every crystal magnetic structure previously obtained as an equilibrium configuration is currently a dynamical solution, after the compatibility with Equation (20) has been checked. At the same time, it has to suffer a damping like solution (21), which restores the unperturbed magnetic configuration dominated by the central field. Back-reaction microstructured field becomes negligible after the time

$$\tau = \frac{\rho_0}{k^2 \nu}\,, \tag{31}$$

defined in Equation (14). A direct consequence is that such structures could exist only if the plasma is quasi-ideal, with very low viscosity coefficient; in particular, this shows how the present analysis thwarts the Standard Model [10, 11]. Let us try to use the phenomenological Shakura prescription for the viscosity, i.e.

$$\nu = \alpha \rho_0 v_S H_0\,, \tag{32}$$

where v_S is the sound velocity and α is a model parameter which determines the strength of turbulence enhancement. Remembering the Standard Disc is nearly Keplerian with $\Omega_K \simeq v_S/H_0$, and using Equation (32) in Equation (31), we obtain

$$\tau_{\text{standard}} \simeq \frac{\varepsilon_z^2}{\alpha}\Omega_K^{-1}\,, \tag{33}$$

whose rough estimates (with the common choice of α in the range $0.01 - 0.1$) are of the order of hours for a X-Ray Binary System, absolutely meaningless on the evolution time-scale. This is to prove how the scheme of magnetic crystal microstructures shows itself as an *aut aut* alternative to the Standard Model, since those kind of configurations are not held by the inbuilt assumptions of the α paradigm, at least if we seek for a quasi-stationary picture on the time-scale of stellar evolution.

On the other hand, relation (31) provides us with constraints on the kind of systems in which those structures could be found. To this purpose, we use Equation (13) and the formula for microscopical resistivity:

$$\eta = \frac{m_e \nu_{ie}}{n_e e^2} \simeq 1.435 \cdot 10^{-8} \frac{\log \Lambda}{\left(T\,[\mathrm{K}]\right)^{3/2}}\ [\mathrm{s}]\,, \tag{34}$$

where the Coulomb Logarithm can be approximated by $\log \Lambda \simeq 20$ at the typical densities of interest. The mean lifetime τ can then be expressed as a function of only the temperature T and the back-reaction length scale $\lambda \doteq 2\pi k^{-1}$.

In Fig. (2) we show $\tau(\lambda)$ for some different values of temperature. The horizontal thick line marks the estimated lifetime (10^7 years) for an accretion disc in High Mass X-Ray Binary Systems, when the disc can continuously replenish matter from the companion star. Microstructures' lifetime has to exceed this value in order to speak of a quasi-stationary configuration on the scale of stellar evolution, but you can see that this threshold is achieved only with very high temperatures: a reasonable value of $10^7\,$K yields the length scale over $10^9\,$cm, which is comparable to the disc size; reducing this scale to a more meaningful $10^7\,$cm needs the temperature to exceed $10^9\,$K, far beyond the typical range. We deduce this model is inconsistent with those time-scales and should not be applied in this way. We point out now that in [4] an estimation of λ in the case of a global structure is provided, and it is of the order of $10^3 - 10^4\,$cm, which gives rise to a negligible lifetime regardless of temperature.

VI. CONCLUSIONS

We discussed how the equilibrium configurations proposed by [1] and following works have to be generalized by means of non-ideal terms (an arbitrarily small but non-zero resistivity is required by kinetic theory, and its effects are not negligible), and how these equilibria evolve in time. The peculiar morphology of magnetic microstructures is preserved, but it has been shown how the dissipative coefficients forbids its maintaining: in particular, the macroscopical viscosity of Shakura Standard Disc destroys it in a few hours or less, and the microscopical plasma resistivity is nevertheless unable to reach the time-scales of stellar evolution in binary systems. In spite of this, we noted that ranges of length and temperature exist (shaded region in Fig. (2)) which are consistent with an evolution in terms of transient phenomena whose lifetime is between months and years. Such a framework could involve plasma discs in the so called Cataclysmic Variables (e.g., DQ Herculis systems: see [14]), which are known to exhibit short-period events, as well as in Black Hole X-Ray Transient systems, which undergo very long periods of quiescence interrupted by non-periodic small-duration times of strong activity – for instance, see the atypical X-ray source GRO J1655-40 [15]. Temperatures in $10^3 - 10^5$ K and length scales in $10^6 - 10^8$ cm are suitable for this kind of objects, suggesting the formation and disruption of magnetic microstructures; the same range is available for the description of XRB systems in the case of no replenishing of matter, since isolated discs have similar lifetimes.

[1] B. Coppi, Phys. Plasmas **12**, 7302 (2005).
[2] B. Coppi and F. Rousseau, Astrophys. J. **641**, 458 (2006).
[3] M. Lattanzi and G. Montani, Europhys. Lett. **89**, 39001 (2010).
[4] G. Montani and R. Benini, Phys. Rev. E **84**, 026406 (2011).
[5] G. S. Bisnovatyi-Kogan and R. V. E. Lovelace, NewAR **45**, 663 (2001).
[6] S. A. Balbus, Ann. Rev. Astron. Astr. **41**, 555 (2003).
[7] V. C. A. Ferraro, MNRAS **97**, 458 (1937).
[8] G. Montani and R. Benini, Gen. Rel. Grav. **43**, 1121 (2011).
[9] R. Benini, G. Montani, and J. Petitta, Europhys. Lett. **96**, 19002 (2011).
[10] N. I. Shakura, Sov. Astron. **16**, 756 (1973).
[11] N. I. Shakura and R. A. Sunyaev, A&A **24**, 337 (1973).
[12] G. Bodo, F. Cattaneo, A. Ferrari, A. Mignone, and P. Rossi, Astrophys. J. **739**, 82 (2011).
[13] C. Zanni, A. Ferrari, R. Rosner, G. Bodo, and S. Massaglia, A&A **469**, 811 (2007).
[14] J. Patterson, PASP **106**, 209 (1994).
[15] A. A. Esin, J.-P. Lasota, and R. I. Hynes, A&A **354**, 987 (2000).